anti-ZSCAN29-RAB-C10



Applications

Competition ELISA	Western Blot	SPR	Spiked IP	Immunofluorescence	IP-MS	ChIP
Pass			Pass	Pass		

^{*}rAb has been tested for the following applications. See below for the experimental details.

Antibody information

rAb ID: anti-ZSCAN29-RAB-C10

Description: recombinant Fab fragment obtained by recombinant antibody (rAb) phage display recognizing *ZSCAN29* protein under non-denaturing conditions; specificity and affinity tested.

Binder type: rAb **Isotype:** IgG1 **Species:** *Homo sapiens* **Produced in:** *E. coli* **rAb tags:** no tag; no tag **Specificity:** reacts with *Homo sapiens* ZSCAN29 **Epitope:** binds to folded domain amino acids 9-104

Storage conditions: short term – store at n 4°C (over 6 months), long term - PBS -20°C or -80°C

Link: http://recombinant-antibodies.org/binders/anti-ZSCAN29-RAB-C10

Antigen information

Protein Name: Zinc finger and SCAN domain-containing protein 29 **HGNC Symbol:** ZSCAN29 **HGNC ID:** 26673 **Species:** *Homo sapiens*

UniProt AC: Q8IWY8 UniProt KB: ZSC29 HUMAN

Protein Sequence:


MSGLNDIFEAQKIEWHEHHHHHHENLYFQSHMENGTNSETFRQRFRRFHYQEVAGPREAFSQLWELCCRWLRPEVRTKE QIVELLVLEQFLTVLPGEIQNWVQEQCPENGEEAVTLVEDLEREPGRPR

 Tag N-terminus:
 MSGLNDIFEAQKIEWHEHHHHHHHHHENLYFQSHM
 Tag C-terminus:

Vector Type: pET15Avi6HT_NESG Vector Link: http://beta.labgeni.us/registries/DNASU/pET15Avi6HT_NESG/
Protein Sequence Position: 9-104 Antigen source: *E. coli* Source Lab: Rutgers Source Lab ID: HR8429A.001


Description: affinity purified recombinant protein

Validation data



Single point competition phage ELISA Plot represents specific binding of a target to the rAb-phage in solution (right bar) in comparison to binding to the target immobilized on the plate surface (left bar). Experimental conditions were calibrated to capture binders with dissociation Constant (K_D): 20nM or lower.

Experimental Conditions: Culture supernatants containing rAbphage were diluted five-fold in phosphate-buffered saline, 0.5% (w/v) BSA, 0.1% (v/v) Tween 20 either with or without soluble antigen competitor at 20 nM. After 1 h incubation at room temperature, the mixtures were transferred to neutravidin coated plates preloaded with 50 µL of 20 nM biotinylated antigen and incubated for 15 min. The plates were washed with phosphatebuffered saline, 0.05% (v/v) Tween 20 and incubated for 30 min with horse radish peroxidase/anti-M13 antibody conjugate (1:5000 dilution). The plates were washed, developed with 3,3',5,5'-Tetramethyl-benzidine/H₂O₂ peroxidase substrate (Thermo Scientific), quenched with 1M H₃PO₄, and the absorbance at 450 nm (A450) was determined.



Spiked IP Tritration curve of rAb against antigen of interest. The K_D values were obtained by the least-squares fitting of fluorescence saturation data.

Dissociation Constant (K_D): < 1 nM

Experimental Conditions:

Spiked IP: Antigen was immobilized to M280 Dynabeads. A rAb, 50 nM, was pulled down from high salt AFC buffer with or without HEK293 lysate (OD280 ~10). Beads were washed with low salt AFC buffer, and the captured antibody was quantified with an anti-Fab fluorophore labeled antibody on a flow cytometer.

Affinity Measurement: Antigen was immobilized to M280 Dynabeads and incubated with a rAb, in varying concentration (100 nM down to 1 nM in three-fold dilutions). Beads were washed with BSET/BSA and quantified.

Buffers:

High salt AFC buffer: 10 mM Tris-HCl, pH 7.9, 420 mM NaCl, 0.1% NP-40

Low salt AFC buffer: 10 mM Tris-HCl, pH 7.9, 100 mM NaCl, 0.1% NP-40

PBSE/BSA: 20 mM Na2HPO4, pH 7.5, 150 mM NaCl, 1 mM

EDTA, 0.5% BSA

PBSET/BSA: PBSE/BSA + 0.1% Tween-20

Immunofluorescence:
Status: Pass
Experimental Conditions: http://recombinant-antibodies.org/protocols/immunofluorescence
IP-MS – immunoprecipitation for mass spectrometric analysis:
Status:
Experimental Conditions: http://recombinant-antibodies.org/protocols/IP-MS
ChIP – chromatin immunoprecipitation:
Status:
Experimental Conditions: Pending
IP – immunoprecipitation:
Status:
Experimental Conditions: Pending
SP Elisa:
Status:
Experimental Conditions: http://recombinant-antibodies.org/protocols/ELISA-IC50-EC50-direct-coating

Visit us at http://recombinant-antibodies.org/

Contact:

Recombinant Antibody Network

admin@recombinant-antibodies.org

The University of Chicago

Knapp Center for Biomedical Discovery Rm. 3240G 900 E. 57th St., Chicago, IL 60637

Phone: +1 (773) 834-2776

University of California, San Franciso

Byers Hall Rm. 503

1700 4th St., San Francisco, CA 94158

Phone: +1 (530) 341-2371

University of Toronto

Best Institute Rm. 117

112 College Avenue, Toronto, Ontario M5G 1L6

Phone: +1 (416) 978-1594