anti-KDM2A-RAB-C324



Applications

Competition ELISA	Western Blot	SPR	Spiked IP	Immunofluorescence	IP-MS	ChIP
Pass				Pass	Pass	

^{*}rAb has been tested for the following applications. See below for the experimental details.

Antibody information

rAb ID: anti-KDM2A-RAB-C324

Description: recombinant Fab fragment obtained by recombinant antibody (rAb) phage display recognizing *KDM2A* protein under non-denaturing conditions; specificity and affinity tested.

Specificity: reacts with *Homo sapiens* KDM2A **Epitope:** binds to folded domain amino acids 1-517 **Storage conditions:** short term – store at n 4°C (over 6 months), long term - PBS -20°C or -80°C

Link: http://recombinant-antibodies.org/binders/anti-KDM2A-RAB-C324

Antigen information

Protein Name: Lysine-specific demethylase 2A

HGNC Symbol: KDM2A HGNC ID: 13606 Species: Homo sapiens

UniProt AC: Q9Y2K7 UniProt KB: KDM2A HUMAN

Protein Sequence:


MSGLNDIFEAQKIEWHEGSAGGSGMEPEEERIRYSQRLRGTMRRRYEDDGISDDEIEGKRTFDLEEKLHTNKYNANFVTF
MEGKDFNVEYIQRGGLRDPLIFKNSDGLGIKMPDPDFTVNDVKMCVGSRRMVDVMDVNTQKGIEMTMAQWTRYYETPEE
EREKLYNVISLEFSHTRLENMVQRPSTVDFIDWVDNMWPRHLKESQTESTNAILEMQYPKVQKYCLMSVRGCYTDFHVDF
GGTSVWYHIHQGGKVFWLIPPTAHNLELYENWLLSGKQGDIFLGDRVSDCQRIELKQGYTFVIPSGWIHAVYTPTDTLVFG
GNFLHSFNIPMQLKIYNIEDRTRVPNKFRYPFYYEMCWYVLERYVYCITNRSHLTKEFQKESLSMDLELNGLESGNGDEEA
VDREPRRLSSRRSVLTSPVANGVNLDYDGLGKTCRSLPSLKKTLAGDSSSDCSRGSHNGQVWDPQCAPRKDRQVHLTHF
ELEGLRCLVDKLESLPLHKKCVPTGIEDEDALIADVKILLEELANSDPKLALTGVPIVQWPGGSGHHHHHH

Tag N-terminus: MSGLNDIFEAQKIEWHEGSAGGSG Tag C-terminus: GGSGHHHHHH

Vector Type: p28BIOH-LIC Vector Link: http://www.thesgc.org/sites/default/files/toronto_vectors/p28BIOH-LIC.pdf Protein Sequence Position: 1-517 Antigen source: *E. coli* Source Lab: SGC Source Lab ID: FBXL11A-A002

Description: affinity purified recombinant protein

Validation data



Single point competition phage ELISA Plot represents specific binding of a target to the rAb-phage in solution (right bar) in comparison to binding to the target immobilized on the plate surface (left bar). Experimental conditions were calibrated to capture binders with dissociation Constant (K_D): 20nM or lower.

Experimental Conditions: Culture supernatants containing rAbphage were diluted five-fold in phosphate-buffered saline, 0.5% (w/v) BSA, 0.1% (v/v) Tween 20 either with or without soluble antigen competitor at 20 nM. After 1 h incubation at room temperature, the mixtures were transferred to neutravidin coated plates preloaded with 50 µL of 20 nM biotinylated antigen and incubated for 15 min. The plates were washed with phosphatebuffered saline, 0.05% (v/v) Tween 20 and incubated for 30 min with horse radish peroxidase/anti-M13 antibody conjugate (1:5000 dilution). The plates were washed, developed with 3,3',5,5'-Tetramethyl-benzidine/H₂O₂ peroxidase substrate (Thermo Scientific), quenched with 1M H₃PO₄, and the absorbance at 450 nm (A450) was determined.

Spiked IP: Status:

Experimental Conditions: Pending

Immunofluorescence:

Status: Pass

Experimental Conditions: https://recombinant-antibodies.org/protocols/#uagb-tabs tab1

IP-MS – immunoprecipitation for mass spectrometric analysis:

Status: Pass

Experimental Conditions: https://recombinant-antibodies.org/protocols/#uagb-tabs tab1

ChIP – chromatin immunoprecipitation:

Status:

Experimental Conditions: Pending

IP - immunoprecipitation:

Status:

Experimental Conditions: Pending

SP Elisa:

Status:

Experimental Conditions: https://recombinant-antibodies.org/protocols/#uagb-tabs tab1

Visit us at http://recombinant-antibodies.org/

Contact:

Recombinant Antibody Network

admin@recombinant-antibodies.org

The University of Chicago

Knapp Center for Biomedical Discovery Rm. 3240G 900 E. 57th St., Chicago, IL 60637

University of California, San Franciso

UCSF Byers Hall Rm. 504 1700 4th St., San Francisco, CA 94158

University of Waterloo

School of Pharmacy 10 A Victoria St. S., Kitchener, Ontario, Canada N2G 1C5