anti-SMU1-RAB-C513



Applications

Competition ELISA	Western Blot	SPR	Spiked IP	Immunofluorescence	IP-MS	ChIP
Pass			Pass	Pass		

^{*}rAb has been tested for the following applications. See below for the experimental details.

Antibody information

rAb ID: anti-SMU1-RAB-C513

Description: recombinant Fab fragment obtained by recombinant antibody (rAb) phage display recognizing *SMU1* protein under non-denaturing conditions; specificity and affinity tested.

Binder type: rAb Isotype: IgG1 Species: Homo sapiens Produced in: E. coli rAb tags: Avi-tag; no tag

Specificity: reacts with *Homo sapiens* SMU1 **Epitope:** binds to folded domain amino acids 1-513 **Storage conditions:** short term – store at n 4°C (over 6 months), long term - PBS -20°C or -80°C

Link: http://recombinant-antibodies.org/binders/anti-SMU1-RAB-C513

Antigen information

Protein Name: Smu-1 suppressor of mec-8 and unc-52 homolog **HGNC Symbol:** SMU1 **HGNC ID:** 18247 **Species:** *Homo sapiens*

UniProt AC: Q2TAY7 UniProt KB: SMU1 HUMAN

Protein Sequence:

MSGLNDIFEAQKIEWHEGSAGGSGMSIEIESSDVIRLIMQYLKENSLHRALATLQEETTVSLNTVDSIESFVADINSGHWDTV LQAIQSLKLPDKTLIDLYEQVVLELIELRELGAARSLLRQTDPMIMLKQTQPERYIHLENLLARSYFDPREAYPDGSSKEKRR AAIAQALAGEVSVVPPSRLMALLGQALKWQQHQGLLPPGMTIDLFRGKAAVKDVEEEKFPTQLSRHIKFGQKSHVECARF SPDGQYLVTGSVDGFIEVWNFTTGKIRKDLKYQAQDNFMMMDDAVLCMCFSRDTEMLATGAQDGKIKVWKIQSGQCLRR FERAHSKGVTCLSFSKDSSQILSASFDQTIRIHGLKSGKTLKEFRGHSSFVNEATFTQDGHYIISASSDGTVKIWNMKTTECS NTFKSLGSTAGTDITVNSVILLPKNPEHFVVCNRSNTVVIMNMQGQIVRSFSSGKREGGDFVCCALSPRGEWIYCVGEDFV LYCFSTVTGKLERTLTVHEKDVIGIAHHPHQNLIATYSEDGLLKLWKPGGSGHHHHHH


Tag N-terminus: MSGLNDIFEAQKIEWHEGSAGGSG Tag C-terminus: GGSGHHHHHHH

Vector Type: p28BIOH-LIC Vector Link: http://www.thesgc.org/sites/default/files/toronto_vectors/p28BIOH-LIC.pdf

Protein Sequence Position: 1-513 Antigen source: E. coli Source Lab: SGC Source Lab ID: SMU1-A001

Description: affinity purified recombinant protein

Validation data



Single point competition phage ELISA Plot represents specific binding of a target to the rAb-phage in solution (right bar) in comparison to binding to the target immobilized on the plate surface (left bar). Experimental conditions were calibrated to capture binders with dissociation Constant (K_D): 20nM or lower.

Experimental Conditions: Culture supernatants containing rAbphage were diluted five-fold in phosphate-buffered saline, 0.5% (w/v) BSA, 0.1% (v/v) Tween 20 either with or without soluble antigen competitor at 20 nM. After 1 h incubation at room temperature, the mixtures were transferred to neutravidin coated plates preloaded with 50 µL of 20 nM biotinylated antigen and incubated for 15 min. The plates were washed with phosphatebuffered saline, 0.05% (v/v) Tween 20 and incubated for 30 min with horse radish peroxidase/anti-M13 antibody conjugate (1:5000 dilution). The plates were washed, developed with peroxidase 3,3',5,5'-Tetramethyl-benzidine/H₂O₂ substrate (Thermo Scientific), quenched with 1M H₃PO₄, and the absorbance at 450 nm (A450) was determined.

Spiked IP: Status: Pass

Experimental Conditions: http://recombinant-antibodies.org/protocols/spiked-IP

Immunofluorescence:

Status: Pass

Experimental Conditions: http://recombinant-antibodies.org/protocols/immunofluorescence

IP-MS – immunoprecipitation for mass spectrometric analysis:

Status:

Experimental Conditions: http://recombinant-antibodies.org/protocols/IP-MS

ChIP – chromatin immunoprecipitation:

Status:

Experimental Conditions: Pending

IP - immunoprecipitation:

Status:

Experimental Conditions: Pending

SP Elisa:

Status:

Experimental Conditions: http://recombinant-antibodies.org/protocols/ELISA-IC50-EC50-direct-coating

Visit us at http://recombinant-antibodies.org/

Contact:

Recombinant Antibody Network

admin@recombinant-antibodies.org

The University of Chicago

Knapp Center for Biomedical Discovery Rm. 3240G 900 E. 57th St., Chicago, IL 60637

Phone: +1 (773) 834-2776

University of California, San Franciso

Byers Hall Rm. 503

1700 4th St., San Francisco, CA 94158

Phone: +1 (530) 341-2371

University of Toronto

Best Institute Rm. 117

112 College Avenue, Toronto, Ontario M5G 1L6

Phone: +1 (416) 978-1594